

## Drug Facts (continued)

Adults and adolescents (12 years and older)	Take 10 drops 3 times daily or as recommended by your healthcare practitioner.
---	--

Children (under 12 years)	Take under the direction of your healthcare practitioner.
---------------------------	---

### Other information

- Do not use if seal is missing or broken.
- Store in a cool, dry place.
- Contents may not fill package in order to accommodate required labeling. Please rely on stated quantity.

### Inactive ingredients

Ethanol (beet), purified water, glycerin

Seroyal 

Made in Belgium for  
Seroyal USA  
Pittsburgh, PA 15275  
1-888-737-6925

V2

UNDA® and design is used under license from Unda S.A.



NDC 62106-1200-0



## CARDUUS PLEX

Homeopathic Preparation

For the temporary relief of symptoms associated with poor digestion.



Contains 45% Alcohol

30 mL (1 fl oz)

## Drug Facts

### Active ingredients . Purpose

Each drop contains:

Belladonna (Deadly nightshade)

Whole Plant 4X

(<1 x 10<sup>-6</sup>% alkaloids, calculated) . . . . .

Stomach pains

Carduus marianus

(Milk thistle) Fruit 2X . . . . .

Cramps

Chelidonium majus

(Greater celandine) Rhizome and Root 2X . . . . .

Indigestion

Cholesterinum (Cholesterol) 4X . . . . .

Abdominal pain

Cinchona succirubra

(Peruvian bark) Bark 3X . . . . .

Flatulence

Colocynthis (Bitter apple)

Fruit Pulp 6X . . . . .

Stomach pain

Leptandra virginica

(Culver's root) Underground

Parts 1X . . . . .

Indigestion

Lycopodium clavatum

(Clubmoss) Spores 4X . . . . .

Bloating

Podophyllum peltatum (Mayapple)

Underground Parts 4X . . . . .

Heartburn

Taraxacum officinale

(Common dandelion)

Whole Plant 2X . . . . .

Hiccups

X is a homeopathic dilution:

see [www.seroyal.com/homeopathics](http://www.seroyal.com/homeopathics)

for details. . . . . →

## Drug Facts (continued)

### Uses

- Helps promote and maintain overall digestive health.
- For the temporary relief of symptoms associated with poor digestion.

Product 'Uses' are based on Homeopathic Materia Medica and are not supported by clinical trials. These 'Uses' have not been evaluated by the Food and Drug Administration.

### Warnings

**Stop use and ask a doctor if symptoms persist or worsen.**

**If pregnant or breastfeeding, ask a health professional before use. Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

### Directions

- To be taken 10 minutes away from food.
- Invert bottle and shake lightly allowing drops to fall directly under the tongue or into ¼ oz of water.
- Hold contents for about 20 seconds and swallow.
- To promote drop flow, invert bottle and shake before each use. →