## Drug Facts (continued)

Adults and adolescents (12 years and older) Take 10 drops three times daily or as recommended by your healthcare practitioner.

Children (under 12 years)

Take under the direction of your healthcare practitioner.

### Other information

- Do not use if seal is missing or broken.
- Store in a cool, dry place.
- Contents may not fill package in order to accommodate required labeling.
   Please rely on stated quantity.

### Inactive ingredients

Ethanol (beet), glycerin, purified water

## Seroyal 🦓

Made in Belgium by UNDA Dist. by Seroyal USA Pittsburgh, PA 15275 1-888-737-6925

BARCODE LOCATION

V2

NDC 62106-1195-0
UNDA

# LOBELIA PLEX Homeopathic Preparation

Assists with the temporary relief of cough.



Contains 76% Alcohol

1 fl oz (30 mL)

## Drug Facts

## Active ingredients Purpose

Each drop contains:
Antimonium sulphuratum aureum
(Golden antimony sulfide) 8X Cougl
Arsenicum iodatum
(Arsenic triiodide) 8X Cougl
Blatta orientalis (Indian cockroach)
Live Insect 6X Cougl
Eriodictyon californicum
(Yerba santa) Aerial Parts 3X Congestion
lpecacuanha (lpecac)
Underground Parts 4X Coug
Lobelia inflata (Indian tobacco)
Whole Plant 8X Cougl
Natrum sulphuricum
(Sodium sulfate) 10X Coug
Sambucus nigra (European elder)
Leaves and Inflorescences 1X Coug
Senega officinalis
(Seneca snakeroot) Root 1X Shortnes

X is a homeopathic dilution: see www.seroyal.com/homeopathics for details.

#### Uses

Assists with the temporary relief of cough.

Product 'Uses' are based on Homeopathic Materia Medica and are not supported by clinical trials. These 'Uses' have not been evaluated by the Food and Drug Administration.

## Drug Facts (continued)

## Warnings

Ask a doctor before use if you have

- Persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema.
- Cough that is accompanied by excessive phlegm (mucus) or fever.

#### Stop use and ask a doctor if

- Cough persists for more than 1 week, tends to recur, or is accompanied by a fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.
   Symptoms persist or worsen.
- Symptoms persist or worsen

If pregnant or breastfeeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

#### Directions

of breath

- To be taken ten minutes away from food.

   Invert bottle and shake lightly allowing.
- Invert bottle and shake lightly allowing drops to fall directly under the tongue or into ¼ oz of water.
- Hold contents for about 20 seconds and swallow.
- To promote drop flow, invert bottle and shake before each use. →