

Metabolic Protein Chocolate

Delicious chocolate-flavored pea protein, nutrient and berry extract formulation

- Combines pea protein, alpha-lipoic acid, chromium, vitamins and berry extracts
- Promotes healthy nutrient and glucose metabolism[‡]
- Provides 15 g of pea protein and 20 mg of berry polyphenols per dose

Metabolic Protein Chocolate helps maintain healthy metabolism with a blend of pea protein, alpha-lipoic acid, chromium, vitamins and polyphenolrich berry extracts. This plant-based protein formula helps build and maintain body tissue, and assists in the building of lean muscle mass when combined with regular resistance training and a healthy balanced diet. Chromium is included as an essential mineral that helps maintain healthy blood glucose levels within the normal range. Vitamin B_{12} plays important roles in red blood cell formation and energy metabolism, while vitamin D is well-recognized for its ability to support immune function and help build strong bones and teeth. This formula also combines alpha-lipoic acid with standardized blueberry and strawberry extracts for their valuable effects on antioxidant defense. Metabolic Protein Chocolate is ideal for individuals looking to increase their intake of vegetarian protein, vitamins, antioxidants and complementary ingredients in a delicious, chocolate-flavored formula.[‡]

SUPPLEMENT FACTS Serving Size 2 Scoops (30.2 g) Servings per Container about 14 AMOUNT PER SERVING % DV Calories 110 3%^ Total Fat 2.5 g 2%^ Total Carbohydrate 5 g 3 g **Dietary Fiber** 11%^ Protein 15 g 30%^ Vitamin D (as cholecalciferol) 10 mcg (400 IU) 50% 440 mcg Vitamin B₁₂ (as methylcobalamin) 18.333% Iron √ 6 mg 37% 200 mcg 571% Chromium (as chromium nicotinate glycinate) Sodium √ 230 mg 10% Potassium V 240 mg 5% 15 g Pea Protein (from pea seed) 300 mg * **DL-Alpha-Lipoic Acid** 25 mg * Curcumin (from turmeric rhizome) -)(Lecithin (from sunflower seed) 600 mg * Strawberry (Fragaria x ananassa) Fruit Extract (25:1) 500 mg (12.5 g Fresh Equivalent) Blueberry (Vaccinium angustifolium) Fruit Std. Extract (30:1) 500 mg * (4% Polyphenols/15 g Fresh Equivalent) * Daily value (DV) not established ^ Percent daily values (DV) are based on a 2,000 calorie diet V Naturally occuring

Other ingredients: Cocoa powder, guar gum, natural chocolate, cream and vanilla cream flavors, rebaudioside A (stevia leaf extract), thaumatin

Recommended Dose

In a glass, add 8 ounces (250 mL) of water to 2 scoops of Metabolic Protein Chocolate and mix. Take once daily or as recommended by your health professional.



Product Code 06454

GenestraBrands.com | 1.888.737.6925

© 2019 Seroyal. All rights reserved.

Metabolic Protein Chocolate

Scientific Rationale:

Skeletal muscle is constantly turning over, with new protein replacing older protein lost to degradation.¹ When the rate of muscle protein synthesis is greater than the rate of protein breakdown, there is a gain in muscle mass.¹ In contrast, a net breakdown of muscle protein occurs when protein degradation exceeds protein synthesis.² Protein breakdown is greater than protein synthesis after exercise, resulting in a negative balance of protein turnover.³ Decreased lean muscle mass also occurs naturally during aging, with approximately 1% of total muscle mass lost each year over the age of 30.⁴ In older adults, this decrease in muscle mass is associated with decreased muscle strength and function, as well as reduced quality of life.⁵

Pea protein helps build and maintain body tissues and provides amino acids that play a role in muscle protein synthesis.⁶ Furthermore, when combined with regular resistance training and a healthy, balanced diet, pea protein can help build lean muscle mass.⁶ In one pilot study, supplementation with pea protein for eight weeks led to improvements in muscle thickness similar to those produced by the same dose of whey protein.⁶ As pea protein is not obtained from common allergenic foods, such as milk, soy, wheat or fish, it is ideal for individuals consuming a restricted diet.7‡

Vitamin D is one of the major nutrients involved in bone health.⁸ It plays an essential role in building strong bones and teeth as it helps in the absorption of calcium, a primary structural component of the skeleton.⁸ Vitamin D also helps to stimulate bone mineralization and maturation, while regulating the differentiation of cells present in bone.⁸ Furthermore, vitamin D receptors are present on most immune cells, demonstrating its importance in maintaining immune function.⁹ Individuals at a particular risk of vitamin D inadequacy include those with little sunlight exposure, the elderly, and dark skinned-individuals (due to reduced ability to synthesize vitamin D from sunlight).^{10‡}

B vitamins play an important role in energy metabolism as cofactors for numerous biochemical reactions in the body.¹¹ Vitamin B₁₂ is particularly critical in the metabolism of the amino acid metabolite homocysteine, and has additional roles supporting immune system function and red blood cell formation.⁹ As vitamin B₁₂ is primarily found in animal and dairy products, individuals consuming plant-based diets may benefit from daily supplementation.⁹ Furthermore, adults over 50 may be at an increased risk of low vitamin B₁₂ levels due to a decreased ability to digest food-bound B₁₂ with age.10‡

GenestraBrands.com | 1.888.737.6925

© 2019 Seroyal. All rights reserved.

Chromium can support the normal metabolism of carbohydrates and fats by regulating healthy glucose uptake into the cells, glycogen formation, fatty acid synthesis, and lipid metabolism.¹²⁻¹⁴ Chromium also can support insulin-mediated cell signaling pathways.^{12-14‡}

Free radicals may be generated by environmental sources (such as heavy metals, drugs or ultraviolet rays) or formed during normal metabolic reactions, playing important roles in cell signaling, differentiation and immune cell function.^{15,16} Under normal cellular conditions, antioxidant protection balances free radical production: however, increases in free radical formation or reduced antioxidant activity can result in oxidative stress.¹⁵ In turn, this can damage lipids, protein and DNA, impairing the integrity of the cell membrane as well as normal cell function.¹⁵ Although the body naturally defends against oxidative stress, these processes can become overwhelmed.¹⁷ In addition, the body's ability to protect against and repair damage due to oxidative stress decreases over time.¹⁸ As such, older individuals may experience particular benefits from increasing their antioxidant intake.17,18‡

Alpha-lipoic acid is an antioxidant compound produced endogenously by mitochondria.¹⁹ It plays an important antioxidant role by directly scavenging reactive oxygen species, chelating metals and regenerating other antioxidants, such as glutathione and vitamins C and E.²⁰ As it is both water- and lipid-soluble, alpha-lipoic acid can provide unique antioxidant support in areas of the cell not normally accessed by other antioxidants.^{20‡}

Strawberries are known to contain a high level of vitamin C, along with a range of phenolic compounds including flavonols, anthocyanins and ellagitannins.²¹ Preclinical research suggests that strawberries help reduce oxidative stress, while promoting the activity of antioxidant enzymes.²² **Blueberries** are also well-recognized for their antioxidant effects, resulting primarily from their high anthocyanin content.^{16,17} Animal research has reported that a diet rich in blueberries provided antioxidant support to the liver by promoting hepatic antioxidant enzyme activity and alleviating chemically induced DNA damage.²³ As they can cross the blood-brain barrier, blueberry anthocyanins may also provide specific support to the brain, an organ especially susceptible to oxidative stress due to its high rate of oxygen consumption.^{17,18‡}

- Vincent, JB, Bennett, R. (2007). In JB. Vincent (Ed.), The Nutritional Biochemistry of Chromium (III) (pp. 139-160). Amsterdam, The Netherlands: Elsevier BV.
 Zoidis, E., Seremelis, I, Kontopoulos, N, Danezis, GP. Antioxidants (Basel). 2018; 7(5). pii: E66.
 Peng, C, Wang, X, Chen, J, Jiao, R, Wang, L, et al. Biomed Res Int. 2014; 2014; 831841.
 Kelly, E., Yaya, P. Weber, JT. Molecules. 2017; 23(1). pii: E26.
 Shukitt Hale, B. Gerontology, 2012; 58(6): 518-23.
 Shukitt Hale, B. Goraca, A.2. Oxid Med Cell Longey. 2015; 2015; 313021.
 Skrovankova, S., Sumczynski, D., Micek, J. Jurikova, T, Sochor, J. Int J Mol Sci. 2015; 16(10): 24673-706.
 Oviero Solis, CJ, Cornejo-Manzo, S, Murillo-Ortiz, BD, Guzman-Barrón, MM, Ramirez-Emiliano, J. Gac Med Mex. 2018; 154(1): 80-86.
 Ma, L, Sun, Z, Zeng, Y, Luo, M, Yang, J. Int J Mol Sci. 2018; 19(9). pii: E2785.



⁺ These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.

<sup>REFERENCES
Wolfe, RR, Jint Soc Sports Nutr. 2017; 14: 30.
Wilson, JM, Fitschen, PJ, Campbell, B, Wilson, G, Zanchi, N, et al. Jint Soc Sports Nutr. 2013; 10(1): 6.
Norton, LE, Layman, DK, JNutr. 2006; 136(2): 5335-5375.
Morley, J. Argiles, J. Fvans, W, Bhasin, S, Cella, D, Deutz, N, et al. JAm Med Dir Assoc. 2010; 11: 391-396.
Kim, JS, Wilson, J. Lee, SR. JNutr Biochem. 2010; 21: 1-13.
Banazsek, A. Townsend, JR, Bender, D, Vantrease, WC, Marshall, AC, Johnson, KD. Sports (Basel). 2019; 7(1). pii: E12.
Krefting, J. J Ren Nutr. 2017; 27 (5): e31-e33.
Weichselbaum, E, Buttriss, JL. Nutr Bull. 2014; 39. 9-73.
Combs, GF. (2012). The Vitamins (4th ed.). USA: Elsevier.
American Dietetic Association. J Am Diet Assoc. 2009; 109: 2073-2085.
Panel on Micronutrients, Subcommittees on Upper Reference Levels of Nutrients and of Interpretation and Use of Dietary Reference Intakes and the SC on the SC of DRI. (2001). Washington, DC: National Academies Press.
Hua, Y, Clark, S. Ren, J, Sreejayan, N. J Nutr Biochem. 2012; 23(4): 313-319.
Newsholme, P, Cruzat, V, Ariuso, F, Keane, K. The Journal of Endocrinology, 221(3): R105-R120.</sup>

^{14.} Vincent, JB, Bennett, R. (2007). In JB. Vincent (Ed.), The Nutritional Biochemistry of Chromium (III) (pp. 139-160). Amsterdam, The