

NDC 62106-0631-0

HOMEOPATHIC

PREPARATION To help promote and maintain normal metabolic functions

Contains 31% Alcohol

1 fl oz (30 mL)

Drug Facts Active inevedients

Kali jodatum 6X ...

Lapis albus (Gastein rock)

| Active ingredients | Purpose |
|------------------------------|------------------|
| Each drop contains: | • |
| Calcarea carbonica 8X | Fatigue |
| Calcarea iodata 4X | Thyroid function |
| Ferrum iodatum 6X | Fatigue |
| Fucus vesiculosus | - |
| (Bladderwrack) Thallus 6X7 | Thyroid function |
| Henar sulnhuris calcareum 8X | Chills |

Fatigue Calcium Fluosilicate 6X Thyroid function

Oryctolagus cuniculus (Rabbit) Exhaustion Orvctolagus cuniculus (Rabbit)

D. 180000

Pituitary Gland 10X Growth control

Drug Facts (continued)

Orvctolagus cuniculus (Rabbit) Snleen 10X Weakness Orvctolagus cuniculus (Rabbit)

Thymus Gland 10X Growth control→

Do not use if seal is missing or broken.

BARCODE LOCATION 883196 13810

Peel for more Drug Facts information

Drug Facts (continued) Oryctolagus cuniculus (Rabbit)

Thyroid Gland 10X Spongia tosta (Roasted Sponge) 6X X is a homeopathic dilution; see

www.seroval.com/homeopathics for details. **Uses** Helps promote and maintain normal metabolic

Growth control

Exhaustion

water

functions Product 'Uses' are based on Homeopathic Materia Medica and are not supported by clinical trials. These 'Uses' have not been evaluated by the Food and Drug Administration.

Warnings Stop use and ask a doctor if symptoms persist or worsen.

If pregnant or breastfeeding, ask a health professional before use.

Drug Facts (continued)

get medical help or contact a Poison Control Center right away.

food. ■ Place drops directly under the tongue or into ¼ oz. of water. Adults and adolescents (12 years and older): Take 10 drops two to three times daily or as recommended by your healthcare practitioner. ■ Children (under 12 years):

Inactive ingredients Ethanol (beet), purified

Other information Store in a cool, dry place.

Take under the direction of your healthcare practitioner.

Directions ■ To be taken 20 minutes away from

Keep out of reach of children. In case of overdose.